

The Toxicogenomic Multiverse: Convergent Recruitment of Proteins Into Animal Venoms

Bryan G. Fry,¹ Kim Roelants,^{2,*,**}
Donald E. Champagne,³ Holger Scheib,⁴
Joel D.A. Tyndall,⁵ Glenn F. King,⁶
Timo J. Nevalainen,⁷ Janette A. Norman,^{8,**}
Richard J. Lewis,⁶ Raymond S. Norton,^{9,**}
Camila Renjifo,¹⁰ and
Ricardo C. Rodríguez de la Vega¹¹

¹Department of Biochemistry and Molecular Biology, Bio21 Institute, University of Melbourne, Melbourne 3010 Australia; email: bgf@unimelb.edu.au

Annu. Rev. Genomics Hum. Genet. 2009.
10:483-511

First published online as a Review in Advance on
July 16, 2009

The *Annual Review of Genomics and Human Genetics*
is online at genom.annualreviews.org

This article's doi:
10.1146/annurev.genom.9.081307.164356

Copyright © 2009 by Annual Reviews.
All rights reserved

1527-8204/09/0922-0483\$20.00

*See page 511 for coauthor affiliations.

Key Words

toxin, phylogeny, evolution, convergence

Abstract

Throughout evolution, numerous proteins have been convergently recruited into the venoms of various animals, including centipedes, cephalopods, cone snails, fish, insects (several independent venom systems), platypus, scorpions, shrews, spiders, toxicoferan reptiles (lizards and snakes), and sea anemones. The protein scaffolds utilized convergently have included AVIT/colipase/prokineticin, CAP, chitinase, cystatin, defensins, hyaluronidase, Kunitz, lectin, lipocalin, natriuretic peptide, peptidase S1, phospholipase A₂, sphingomyelinase D, and SPRY. Many of these same venom protein types have also been convergently recruited for use in the hematophagous gland secretions of invertebrates (e.g., fleas, leeches, kissing bugs, mosquitoes, and ticks) and vertebrates (e.g., vampire bats). Here, we discuss a number of overarching structural, functional, and evolutionary generalities of the protein families from which these toxins have been frequently recruited and propose a revised and expanded working definition for venom. Given the large number of striking similarities between the protein compositions of conventional venoms and hematophagous secretions, we argue that the latter should also fall under the same definition.

INTRODUCTION

Venom systems are key evolutionary innovations in a broad phylogenetic range of animal lineages and are used for defense, competitor deterrence, or predation (or a combination thereof). Apart from the extensively studied, medically important clades (snakes, scorpions, and spiders), venomous animals include sea anemones, jellyfish, sea snails, cephalopods, centipedes, several insect orders, echinoderms, fish, lizards, and even some mammals (platypus and shrews). Modes of venom delivery are likewise diverse and include barbs, beaks, fangs or modified teeth, harpoons, nematocysts, pinchers, proboscises, spines, sprays, spurs, and stingers. Targets of venom action include virtually all major physiological pathways and tissue types accessible by the bloodstream.

Venom delivery systems sometimes show great variation in structure and complexity, even within a single venomous clade. Within the reptile clade *Toxicofera*, for example, venom was a single ancient innovation; iguanian and anguimorph lizards share a common venomous ancestor with snakes (56). In iguanian species, venom glands develop only to an incipient stage and seem to have little ecological relevance. However, anguimorph lizards (e.g., monitor lizards and Gila monsters) have a derived, encapsulated, and compartmentalized mandibular venom gland with developmentally suppressed maxillary glands (56). In contrast, snakes possess highly developed but non-compartmentalized maxillary venom glands with developmentally suppressed mandibular glands. Spread across the advanced snakes is an enormous variation of fang and venom gland morphologies (55) that arose from the ancient serpent dental uncoupling (156). Further, within the advanced snakes, divergence has resulted in four derived lineages [twice within the *Lamprophiidae* family (*Atractaspis* and *Homoroselaps* genera), and once each at the base of the *Elapidae* and *Viperidae* families] that possess intricate hollow-front, fanged, and high-pressure delivery systems (55).

Venoms across the Kingdom *Animalia* are complex mixtures that include variable

combinations of proteins (ranging from multi-unit globular enzymes to small peptides), salts, and organic molecules such as polyamines, amino acids, and neurotransmitters (49, 54, 72, 117, 144). Proteins found in venoms are the result of toxin recruitment events in which an ordinary protein gene, typically one involved in a key regulatory process, is duplicated, and the new gene is selectively expressed in the venom gland. In many cases, such toxin genes were amplified to obtain multigene families with extensive neofunctionalization (54, 81), followed by the deletion of some copies and degradation of others to nonfunctional copies or pseudogenes (59). The newly created toxin multigene families often preserve the molecular scaffold (including the tertiary structure) of the ancestral protein, but key functional residues outside the core scaffold are modified to acquire a myriad of newly derived activities (38, 53, 54, 59, 123).

Despite the extraordinary diversity in the structure and function of animal venom systems, several protein groups have been convergently recruited for use as venom toxins in multiple animal lineages (**Table 1**). The convergent origin of toxins across the entire metazoan spectrum suggests that there are functional and/or structural constraints on the evolution of animal venoms. Intriguingly, some of the proteins recruited as venom toxins are also recruited by hematophagous insects for utilization in their feeding secretions; with a wide range of convergent activities within the neurological and hematological systems. The aim of this review is thus twofold: (*a*) to consider the structural and functional constraints on recruitment and evolution of venom proteins and (*b*) to explore the implications of these findings for the definition of what constitutes a venomous animal.

CONVERGENTLY RECRUITED PROTEIN FAMILIES

AVIT/Colipase/Prokineticin

First isolated from black mamba venom (mamba intestinal toxin 1, MIT-1) (21, 142), AVIT peptides have subsequently been

Table 1 Convergently recruited proteins*

	AVIT	CAP	Chi	Cys	Def	Hya	Kun	Lec	Lip	Nat	PS1	PLA ₂	Sm-D	SPRY
Amphibian	X													
Cephalopod		X	X			X					X	X		
Cnidarian							X					X		
Cone snail		X					X							
Fish						X		X						X
Insect (bristle)				X				X	X		X	X		
Insect (proboscis)		X		X		X	X	X	X		X	X		
Insect (stinger)		X	X			X	X				X	X		
Platypus					X					X				
Scorpion		X			X	X	X					X		
Shrew											X			
Spider	X	X				X	X(2)						X	
Reptile	X	X		X(2)	X	X	X	X	X	X	X	X(3)		X
Tick	X	X		X			X		X		X	X	X	

*Abbreviations used: CAP, CRISP (cysteine rich secretory proteins), antigen 5 (Ag5) and pathogenesis-related (PR-1) proteins; Chi, chitinase; Cys, cystatin; Def, defensin; Hya, hyaluronidase; Kun, kunitz; lec, lectin; Lip, lipocalin; Nat, natriuretic peptide; PS1, peptidase S1; PLA₂, phospholipase A₂; Sm-D, sphingomyelinase.

identified in the venom of monitor lizards as part of the Toxicofera reptile core chemical arsenal (56). Similar bioactive peptides are found in the defensive skin secretions of *Bombina* fire-bellied toads (Bm8, Bo8, and Bv8) (30, 31, 102). AVIT scaffold peptides found in reptile venom and *Bombina* skin secretion defensive toxins are potent agonists of mammalian prokineticin receptors, and their binding mimics the effect of an endogenous prokineticin overdose. This effect is reinforced by their highly elevated efficiency in binding mammal receptors; both Bv8 and MIT-1 bind to mammal PKR1 and PKR2 (prokineticin receptors 1 and 2) with an affinity that exceeds that of the normal mammalian peptide PK2 (prokineticin 2) by one order of magnitude and that of PK1 (prokineticin 1) by two orders of magnitude (108). Their toxicity is mainly determined by their two potent short-term physiological effects (within minutes), namely gastric smooth muscle contraction and hyperalgesia (increased pain sensitivity). In a longer time frame, Bv8 additionally shows an anorexogenic effect (108), inhibiting the feeding of the affected predator, even in food-deprived situations. Venom of both mygalomorph and araneomorph spider taxa contain

peptides (MIT-like atracotoxins, ACTX) that show a similar motif as vertebrate AVIT peptides consisting of 10 conserved cysteine residues (148, 162). However, these peptides test negative in assays of PK1/PK2 activity, lack the N-terminal AVIT sequence, and are much more diverse in amino acid composition, sequence length, and scaffold shape. Consistent with the likelihood that they fulfill distinct functions, comparative three-dimensional modeling of AVIT and MIT-like ACTX peptides has shown dissimilar tertiary structures and markedly different surface chemistries. These results are perhaps not surprising because insects, which constitute the majority of spider prey, appear to lack the PK/PKR system. So far, insecticidal assays have not revealed a lethal effect, even at high doses (148). Nevertheless, the diversity of MIT-like ACTX peptides within individual species, their presence in a broad phylogenetic range of spiders (suggesting their ancient recruitment and long-term existence in spider venom), and their observed resistance to proteolytic breakdown clearly point to an adaptive role in spider venom. Recently, peptides similar to the spider venom peptides were also found in the hematophagous secretion

CAP: CRISP (cysteine-rich secretory proteins), antigen 5 (Ag5), and pathogenesis-related (PR-1)

glands of the tick *Ixodes scapularis*, where they potentially play a role in its feeding (128).

CAP [Cysteine-Rich Secretory Proteins (CRISP), Antigen 5 (Ag5), and Pathogenesis-Related (PR-1)] Proteins

Snake venom CRISP scaffold toxins with the C-terminal cysteine-rich domain (CRD) motifs inhibit a number of ion channels, including cyclic nucleotide-gated channels, ryanodine receptor channels, and Ca^{2+} and K^{+} channels (23, 24, 105, 112, 113, 158, 167, 168). Other forms of CAP scaffolds are utilized in the venoms of cephalopods, cone snails, stinging insects, scorpions, and spiders (Figure 1). Tex31, the first of the cone snail proteins to be isolated from the venom duct, possesses proteolytic activity, which has been attributed to the PR-1

domain (100). The cone snail proteins have a C-terminal cysteine-rich domain that is divergent from that found in the venom CRISP proteins, reflecting a different body protein being used in the toxin recruitment event. CAP domain proteins are the dominant allergy-inducing toxins in hymenopteran venoms (47).

Related proteins have been convergently recruited in the feeding secretion of animals from hematophagous taxa in three distinct insect orders: Diptera [mosquitoes (12), sandflies (75), biting midges (26), tsetse flies (86), and muscid flies (4)], Hemiptera [triatomine bugs (129)], and Siphonoptera [fleas (9)]. Within some of these taxa, multiple isoforms are present, contributing to a complex feeding-secretion protein mixture. For example, the transcriptome of *Aedes aegypti* hematophagous secretion glands includes three distinct CAP members (130), and four are found in *Anopheles gambiae* (12). The




Figure 1

Bayesian phylogenetic reconstruction and molecular modeling of representative cysteine-rich secretory proteins (CRISP), antigen 5 (Ag5), and pathogenesis-related (PR-1) (CAP) proteins. Toxic mutant clades are shown in red. In the molecular modeling of representative proteins, blue surface areas indicate positive charges, red indicates negative charges, and model pairs show two sides of the protein rotated by 180°. Methods as per Reference 56.

proteins found in the hematophagus secretions of soft (*Argas*) versus hard (*Ixodes*) ticks represent separate recruitment events (**Figure 1**). The activities of CAP proteins in the insect and tick hematophagous feeding secretions remain to be elucidated. CRISP proteins have also been identified in the buccal glands of the lamprey *Lethenteron japonicum*, a hematophagous parasite of fish. Similar to the reptile toxins, the lamprey buccal secretion CRISPs block L-type voltage-gated Ca^{2+} channels (i.e., Ca_v1), acting as a vasodilator to facilitate the parasite's hematophagy (73).

All arthropod CAP proteins, whether expressed in venom or hematophagous-feeding secretions, differ from those of most other metazoan lineages in lacking a CRD domain. This feature is shared by a CAP protein secreted by the infective nematode *Ancylostoma caninum* during the larval transition to its parasitic stage (67). These worms may lack the CRD domain by common ancestry with arthropods. Both Nematoda and Arthropoda belong to the clade Ecdysozoa, so the loss of the CRD domain may have happened in an early ecdysozoan ancestor rather than in only an arthropod ancestor. According to the conventional structural definition, these Ecdysozoa sequences are not CAP proteins. However, phylogenetic analysis shows they are clearly nested within this protein family (**Figure 1**). The most common CAP/SCP topology does not contain the CRD domain; actually, the CAP-CRISP combination appears to be restricted to the metazoan lineage. Cnidarians are the only clade where a monodomain CRISP (without N-terminal CAP) has been found; therefore, the CAP-CRISP combination appears to be a metazoan innovation. Thus, the biodiversity of CAP proteins is greater than the artificially constrained definition of what constitutes such a protein type, which has focused on a scaffold derivation rather than the molecular ancestral condition.

Chitinase

Chitinase is present in the venom of the braconid wasps *Chelonus* spp. (84), which

endoparasitize the lepidopteran *Trichoplusia ni*, simultaneously ovipositing eggs and injecting venom into the host's eggs. The biological role of this venom chitinase is unclear because chitin appears to be largely absent from both the oviposited wasp eggs and the egg chorion of the host lepidopteran (84). The wasp-venom chitinase has significantly diverged from nonvenom arthropod chitinases ($\leq 45\%$ identity at the amino acid sequence level), and thus it may have acquired new substrate specificity. Chitinase from the posterior salivary gland of the Southern Sand Octopus *Octopus kaurina* (GenBank accession number EU790591) is quite similar in form to the enzyme present in wasp venom. A highly divergent chitinase was also recently discovered in the hematophagous-feeding secretions of soft tick *Argas monolakensis* (91). Birds are the primary host for this tick, and hence the biological role of the hematophagous secretion chitinase is also unknown.

Cystatin

Cystatin-scaffold toxins are found in reptile venoms (56) and also that of the *Lonomia* caterpillar (155), with differing conservation of the characterized functional residues (45). The forms in the tick and mosquito hematophagous secretions also contain mutations in the protease-inhibiting reactive site (63, 128, 130, 173).

Defensins

Defensins are among the most widely distributed innate immunity-related antimicrobial peptides (AMPs) (164, 171). Two structural classes are recognized: the phylogenetically related α -, β -, and θ -defensins that are exclusive to vertebrates (99) and the cysteine-stabilized α/β ($\text{CS}\alpha\beta$) defensins that are found in plants, fungi, nematodes, and mussels as well as in the arthropod classes Insecta and Arachnida (164, 174).

The β -defensin-scaffold toxins characterized from *Crotalus* snake venoms are neurotoxins that modify voltage-gated Na^+ (Na_v)




Figure 2

Bayesian phylogenetic reconstruction of representative defensin proteins. Toxic mutant clades are shown in red. Methods as per Reference 56.

channels, resulting in a potent analgesic effect and significant myotoxicity (151). Because of poor phylogenetic signal, it is unclear whether the form from *Pogona* lizard incipient venom glands represents a separate recruitment event (Figure 2). However, it is quite clear that a unique form in helodermatid lizard venoms, the lethal toxin 1 (LT1) multigene family, represents a separate recruitment event. This unique toxin type is constructed by four tandemly repeated β -defensin motifs, resulting in a single protein with a novel multidomain topology (Fry et al., unpublished results; Genbank accession number EU790964). This shows that high toxin diversity in venoms is attained not only by the expression of multiple paralogs, but also by domain variation in a multidomain single-product gene formed by tandem repeats within an ancestral monodomain gene. The defensin-like peptides isolated from platypus venom are

related to the β -defensin class found only in vertebrates but their bioactivities remain to be elucidated (152).

Remarkably, most scorpion venom neurotoxins adopt the CS $\alpha\beta$ motif, representing an explosive venom-specific gene expansion (124). Although the wide phylogenetic spread of CS $\alpha\beta$ defensins has led some authors to propose a common and very ancient origin of these peptides (53, 175), the high sequence variation prevents reliable phylogenetic reconstructions (39). Nonetheless, bona fide CS $\alpha\beta$ defensin peptides (i.e., those with immune-related function) have been identified in the hematophagous secretion glands of ticks and insects (e.g., 9, 14, 150, 172). Proteins that show remarkable similarity to tick defensins (more than 80% similarity) have been found in scorpion hemolymph (46), supporting a paralogous relationship between CS $\alpha\beta$ defensins and scorpion neurotoxins (37). Indeed, gene-structure conservation and phylogenetic analyses support a paralogous relationship between scorpion defensins and three families of voltage-gated K⁺ (K_V) channel blockers from scorpion venom, likely derived from two independent recruitment events (43).

In the case of mosquitoes, the hematophagous secretion defensins [e.g., defensin A1 (130)] are identical to those produced endogenously as a component of the microbial infection response. A significant volume of hematophagous secretion is reingested with either blood or sugar meals (some hematophagous flies also feed on nectars as supplementary sugar sources), and it has been assumed that these defensins (and other antimicrobial peptides/proteins including cercropins and lysozyme) inhibit bacterial growth in the insect crop or gut. Although this does not preclude a functional toxic role in the vertebrate host, it is suggestive that no hematophagous secretion-specific isoforms (such as may have been produced by selection to fit targets in the vertebrate) have been discovered yet.

Hyaluronidase

Hyaluronidase is found in the venom of toxicoforan reptiles [with isoforms sequences generated in the case of snakes (55, 65) and helodermatid lizards (Fry et al., unpublished results; Genbank accession number EU790961)], stonefish (110), and hymenopterans (61, 77) (**Figure 3**). In addition to venoms for which protein sequences have been obtained, hyaluronidase activity has been reported in octopus (147), spider (107), and scorpion venoms (15), but the corresponding enzymes have not yet been sequenced. Hyaluronidase has also been identified in the hematophagous secretion of dipteran insects (28, 129). Hyaluronidase enzymes across animal taxa show lower levels of sequence diversity than other toxins, and no new activities have been reported for either venom or body forms. Those recruited most likely

act primarily as diffusion factors, enhancing tissue permeability to allow a more efficient spreading of toxins or hemostatic factors (153).

Kunitz-Type Peptides

Kunitz-scaffold toxins have been found in the venoms of snakes (55, 66), sea anemones (10, 17, 69, 101), the solitary wasp *Anopilus samariensis* (68), the scorpion *Hadrurus gertschi* (139), the polychaete worm *Sabellastarte magnifica* (UniProt accession number P84875), the snail *Conus stratius* (16), and the gorgonian coral *Melithaea caledonica* (UniProt accession number P82968). This peptide has also been recruited twice into spider venoms; the scaffold-type in the araneomorph spider *Araneus ventricosus* is phylogenetically distinct from that characterized from the mygalomorph *Ornithoctonus huwena* (**Figure 4**). There is considerable derivation of activity. For example, some snake




Figure 3

Bayesian phylogenetic reconstruction and molecular modeling of representative hyaluronidase enzymes. Toxic mutant clades are shown in red. In the molecular modeling of representative proteins, blue surface areas indicate positive charges, red indicates negative charges, and model pairs show two sides of the protein rotated by 180°. Methods as per Reference 56.




Figure 4

Bayesian phylogenetic and molecular modeling reconstruction of representative Kunitz peptides. In the molecular modeling of representative proteins, blue surface areas indicate positive charges, red indicates negative charges, and model pairs show two sides of the protein rotated by 180°. Toxic mutant clades are shown in red. Methods as per Reference 56.

venom forms inhibit the blood coagulation enzymes plasmin and thrombin (48), whereas others have neurotoxic activity targeting L-type Ca^{2+} channels (141) and K_V channels (66).

Kunitz peptides are also a major constituent of ticks and insect hematophagous secretions, inhibiting the function of blood factor Xa (26, 161). A number of other double-Kunitz domain proteins have also been identified in the hematophagy gland transcriptome of a nematoceran fly, *Culicoides sonorensis* (25). Although the functions of these proteins have not been experimentally demonstrated, three of them (named TFPI1-TFPI3) are presumed to be tissue factor pathway inhibitors on the basis of the pronounced antifactor-Xa activity of hematophagous secretion of this species and their apparent orthology (32% identity and 38% similarity at the amino acid sequence level) with the known antifactor-Xa proteins of tick feeding secretion (52).

Lectin

Lectin-scaffold toxins in snake venoms act via a number of blood coagulation pathways (104) and are also potent anticoagulant toxins in *Lonomia* caterpillar venom (155). In contrast, the lectins in stonefish venom have myotoxic effects (89). C-type lectin proteins have also been found in the feeding secretion of blood-sucking insects, such as *Lutzomyia* sandflies (154) and *Aedes* and *Culex* mosquitoes (130, 131), and may also have an antihemostatic function (11, 130).

Lipocalin

Lonomia obliqua venom contains a number of lipocalin-scaffold variants (134, 155), and one of these, named Lopap, has a unique serine protease-like activity that activates prothrombin (127). There are currently no reports of other lipocalin-scaffold derivations with protease activity.

Several groups of hematophagous arthropods have independently recruited this protein family to serve a variety of antihemostatic functions. Arthropod lipocalins display pronounced molecular biodiversity, such as in the Reduviidae family of predatory insects, which includes the triatomids or kissing bugs (7). For example, 341 scaffold isoforms were recovered from a hematophagous secretion gland cDNA library of *Triatoma brasiliensis* (136). In *Rhodnius prolixus*, the nitrophorins, lipocalin-scaffolds with an unique heme-binding domain, function to store nitric oxide and deliver it to the skin during feeding (5, 27). These scaffold variants also sequester histamine, preventing the host response to histamine released from mast cells and platelets (133), and inhibit coagulation by interfering with factor IXa (64, 132). Another lipocalin (nitrophorin 7, without a heme group) inhibits coagulation by binding to anionic membrane phospholipids, interfering with assembly of procoagulant complexes (8). Yet other lipocalins disrupt hemostasis by sequestering ADP, thus preventing platelet activation and aggregation (50, 51), and by sequestering amines such as serotonin, thereby preventing vasoconstriction and possibly coagulation (6). Species of the bug genus *Triatoma* lack nitrophorins, but diverse lipocalins nevertheless fulfill a variety of antihemostatic roles. Characterized *Triatoma* lipocalins include pallidipin (115), triplatin (103) (which has anticollagen activity), and the antithrombin molecule triabin (114).

Bioactive lipocalin-scaffold variants also independently account for several of the pharmacological activities in the bloodmeal feeding secretion of ticks (93). Moubatin from *Ornithodoros moubata* inhibits collagen-mediated platelet aggregation (160) through a mechanism that includes binding thromboxane A₂ and leukotriene B₄ (90, 96). A second lipocalin from the same tick has 46% amino acid sequence identity with moubatin, but it specifically inhibits the complement pathway by antagonizing activation of complement component C5 and antagonizing C5 activation (97, 116). Two related lipocalins, TGSP4 (from

O. savignyi) and AM-33 (from *Argas monolakensis*), bind cysteinyl leukotrienes with high affinity (95). Cysteinyl leukotrienes, produced by neutrophils, macrophages, basophils, and mast cells in response to injury, increase endothelial permeability leading to edema; this effect would be expected to lead to occlusion of blood vessels and accumulation of a nutritionally deficient serous-rich and erythrocyte-poor fluid at the tick feeding site. Sequestration of these molecules helps ensure a nutritionally adequate meal, rich in erythrocytes (95). FS-HBP (female-specific histamine-binding protein) is a lipocalin expressed specifically by female hard ticks (*Rhipicephalus*) that binds histamine in a manner completely distinct from the *Rhodnius* nitrophorins: Rather than binding through a heme moiety, this protein has two histamine-binding pockets, one low affinity and one high affinity, in the interior of the β -barrel structure (118, 119, 135). Additionally, the lipocalins monomine and monotonin from *O. monolakensis* inhibit vasoconstriction by binding serotonin (5-hydroxytryptamine) as well as histamine (97). These molecules have a single amine-binding pocket and appear to have acquired their amine-binding properties via a separate evolutionary event distinct from the evolution of FS-HBP (97). Analysis of tick hematophagous-feeding secretion gland cDNA libraries has revealed the presence of several other lipocalins of unknown function (128). Lipocalins are also found in the feeding secretion of bloodmeal dipterans, such as the mosquito *Aedes aegypti*, and these forms presumably have anticoagulant activity (130).

Natriuretic Peptides

Natriuretic peptides have been recruited into platypus (41) and reptile (56, 57) venoms and are potent hypotensive toxins. Upstream of the natriuretic peptide-encoding region, the snakes have further evolved either BPP (bradykinin-potentiating peptides) (143) or metalloprotease-inhibiting peptides (157) that are posttranslationally liberated. The helodermatid lizards have independently evolved

PS1: peptidase S1
PLA₂: phospholipase A₂

tandemly repeated proline-rich hypotensive peptides (the helokinestatin peptides) in the same upstream region of the natriuretic peptide gene (Fry et al., unpublished results; Genbank accession number EU790965), and the additional peptides are similarly posttranslationally liberated.

Peptidase S1

The kallikrein-scaffold forms of peptidase S1 (PS1) toxins found in reptile venoms exert a broad variety of activities, ranging from

liberation of kinins from circulating kininogen (ancestral toxic activity) to the cleavage of fibrinogen by a basally derived form (Figure 5). As in the case of the ancestral reptile venom forms, the kallikrein toxins found in *Blarina* shrews also liberate kinins (79, 80). However, the PS1 toxins found in *Lonomia* caterpillar venoms possess fibrinolytic activity (3) (Figure 5), thereby constituting a case of convergent derivation with the reptilian peptidases. The venom of the parasitic wasp *Cotesia rubecula* contains a PS1-scaffold-derived protein (Vn50), which, similarly to the insect PS1 homologs from which




Figure 5

Bayesian phylogenetic reconstruction and molecular modeling of representative peptidase S1 enzymes. Toxic mutant clades are shown in red. In the molecular modeling of representative proteins, blue surface areas indicate positive charges, red indicates negative charges, and model pairs show two sides of the protein rotated by 180°. Methods as per Reference 56.

it evolved, lacks a functional catalytic triad (His-Asp-Ser) because the serine residue is replaced by a glycine (13). The insect PS1 homologs from which Vn50 is derived are involved in a proteolytic cascade that causes hemolymph melanization, which is a primary defense reaction upon infection or invasion by parasites. Vn50 instead inhibits hemolymph melanization, and given its strong structural similarity, it is postulated to be a competitive inhibitor of endogenous peptidase S1 homologs in the host.

PS1 scaffolds in cephalopod posterior salivary glands exhibit a molecular diversity comparable to the diversity observed in the PS1 scaffolds of venomous reptiles (Genbank accession number EU790592–EU790607). Their variation in intraloop (functional) residues appears to constitute an adaptive neofunctionalization pattern that seems analogous to those patterns observed in other multigene toxin families. However, knowledge of the activity of PS1 from cephalopods awaits detailed functional analyses.

The PS1 from the blood-feeding secretion of the vampire bat (*Desmodus rotundus*) is a powerful plasminogen activator (82, 138). PS1s are also widely distributed in the feeding secretions of hematophagous arthropods, including hard ticks (166), nematoceran flies [including mosquitoes (130) and biting midges (26)], and triatomine bugs (136); each of these represents a separate evolutionary recruitment event (Figure 5). Although the specific function of any of these proteins is not known, the presence of CUB domains (named for a class of compounds including complement subcomponents C1r/C1s, Uegf, and bone morphogenic protein-1) suggests substrate specificity (130). These serine proteases are possibly targeted to specific host proteins at the bite site, or they may be involved in activating other proteins following secretion into the host.

Phospholipase A₂

Phospholipase A₂ (PLA₂) enzymes have been convergently recruited into cephalopods,

cnidarians, multiple insect orders (into the toxic arsenals associated with bristles, proboscises, and stingers), arachnids (scorpions, spiders, and ticks), and reptiles (three occasions: twice into the advanced snake venoms and once into anguimorph lizard venom). So far, Group-IA, G-IIA, G-IIB, G-III, G-IX, and G-XII PLA₂ scaffolds have been recruited into venoms (137). The molecular origin and evolutionary relationships have been investigated for G-I, G-II, and G-III PLA₂s (2, 35, 54, 58, 149). Group III PLA₂s are particularly unique in having been recruited independently into four venomous lineages. Types from sea anemone venoms do not associate phylogenetically with any of the currently defined PLA₂ types (109). Derived toxic functions in reptile venoms include antiplatelet, myotoxic, and neurotoxic activities (54, 78). The neurotoxicity of venom PLA₂s depends on their ability to bind to specific target proteins that act as receptors/acceptors in specific organs and tissues (126). After binding to the target, the toxic PLA₂ may induce its effects by mechanisms that are either dependent on or independent of its catalytic activity (78).

Sphingomyelinase D

Sphingomyelinase D (SMaseD) is a magnesium-dependent subclass of toxic enzymes until recently known only from the pathogenic bacterial genus *Corynebacterium* and from the uniquely dermonecrotic venoms in two closely related spider genera (18, 19). Smase D enzymes in both lineages contain similar C-terminal plug motifs and, in the absence of phylogenetic analyses, the disjunct presence of this motif in both lineages was put forward as evidence for a lateral gene transfer event (32). However, SmaseD homologs have recently been identified in ticks (UniProt accession number Q202J4), and BLAST searches reveal the presence of homologous enzymes in other bacteria (*Arcanobacteria* UniProt accession number Q59121) and some fungi (*Aspergillus* and *Coccidioides* UniProt accession numbers Q2UAL9, Q2UKE8, Q2U8X2, and Q1DU31)




Figure 6

Bayesian phylogenetic reconstruction of representative sphingomyelinase enzymes. Toxic mutant clades are shown in red. Methods as per Reference 56.

as well (Figure 6). Phylogenetic analyses show that the spider and tick sequences are closely related and together form a clade that is very distinct from the bacterial and fungal enzymes, which in turn are more similar to each other. A close relationship between tick and spider enzymes does not fit with the originally postulated lateral gene transfer scenario between bacteria and spiders and would push such an event back in time to at least the last common ancestor of both arthropod lineages. If the bacterial forms were the result of lateral gene transfer, then the sequences would be nested within the spider lineages. However, strong structural dissimilarity between the arachnid SmaseD and the bacterial and fungal enzymes (evidenced by large pairwise distances and the long intervening branch in the phylogram of Figure 6) is consistent with a far more ancient divergence between both groups. In addition, a single lateral gene transfer event is not enough to explain the presence of homologs in fungi as well. As an alternative to the lateral gene transfer hypothesis, we postulate that the similarity between bacterial and spider SmaseD arose by convergent evolution in distinct lines of GDPD (glycerophosphodiester phosphodiesterase) enzymes. This scenario gains support from the fact that both the fungal and the tick enzymes

contain C-terminal plug motifs that differ from the identical motifs in the bacterial and spider toxins. In SmaseD of both *Corynebacterium* (bacteria) and *Loxosceles* (spider), Asn residues at position 278 contribute a backbone hydrogen bond, causing subsequent residues to form a type-I β -turn, which blocks the N-terminal end of the TIM (triosephosphate isomerase) barrel (32). In the enzymes of the tick, fungi, and non-*Corynebacterium* bacteria, various other residues occupy position 278 (Asp in the tick; Ala, Lys, Ser, or Thr in fungi; and Ile in *Arcanobacterium*), potentially affecting the plugging of the barrel.

SPRY/Concavalin A-Like Lectins

SPRY toxins have been sequenced in the venoms of toxicoforan reptiles and stonefish. In reptile venoms, they are an ancestral type shared between lizard and snake venoms (56). The bioactivity of the lizard form remains to be elucidated, but the isoform characterized from *Ophiophagus hannah* snake venom produces hypolocomotion and hyperalgesia, perhaps by acting directly on the central nervous system (125). SPRY toxins in stonefish venoms produce lethal hemolysis through pore formation in cell membranes

while inducing potent endothelium-dependent hypotension and also irreversibly interfering with neuromuscular function (29, 88, 146). In addition, these toxins display edema-inducing activity, increase vascular permeability, induce endothelium-dependent vasorelaxation and platelet aggregation, and are myotoxic.

WHAT DETERMINES THE SUITABILITY OF A BODY PROTEIN FOR TOXIN RECRUITMENT?

In contrast to the tremendous diversity of venomous organisms, the number of different venom protein scaffolds appears to be rather restricted. The high proportion of convergently recruited protein families suggests that there are structural and/or functional constraints as to what makes a protein suitable for recruitment as a toxin. Although protein families that have recurrently been recruited span a broad spectrum of different structures and biochemical activities, we notice that many of them share a number of generalities.

First, all known toxins represent secretory proteins. The precursor peptides of all toxins contain an N-terminal signal peptide that is excised to form a functional protein. To date, there are no documented cases of any toxin precursor lacking an N-terminal signal peptide. Even if a nonsecreted body protein might have a potential toxic side effect, its gene would require the addition of a segment encoding a signal peptide either by interlocus gene conversion (nonreciprocal recombination) and retrotransposition or by exon shuffling of a signal peptide protomodule (122). Conversely, the possibility that alternative secretion mechanisms, independent of signal-peptide processing, have evolved in at least some venomous lineages cannot be eliminated. For example, the AMP buforin I, secreted in the stomach of the toad *Bufo gargarizans*, is a derivative of the nucleus-specific protein histone H2A. After proteolytic cleavage from the H2A precursor, it becomes a functional AMP (76). However, the cDNA clone of the H2A

precursor transcript remarkably does not seem to contain an N-terminal signal peptide. Hence, the mechanism that governs the secretion of buforin I is currently unknown. One potential explanation is that buforin I, despite being identical to an N-terminal segment of H2A, is actually encoded by a paralogous gene that obtained a signal peptide after its duplication from the H2A gene, but this remains to be tested. Similar H2A-derived AMPs have been reported from teleost fish (20, 121), indicating that this mechanism may be widespread or evolved convergently in divergent animal lineages. If alternative modes of secretory peptide processing exist for AMPs, they may occasionally have evolved for toxins as well.

Second, protein families from which toxins have been convergently recruited are functionally versatile, but proteins within each family seem to share an underlying biochemistry that has remained remarkably uniform throughout the Kingdom Animalia (and often throughout eukaryotes). Despite evolving very different metabolic, physiological, and immunological processes and having been adapted to the ecological needs of the organisms in which they are expressed, homologs of these proteins often perform the same or very similar fundamental biochemical reactions. For example, the myriad of metabolic functions performed by peptidase S1s are all based on the same underlying biochemical reaction, which is the hydrolysis of specific peptide bonds. Similarly, the biological functions of hyaluronidase enzymes all scale down to the hydrolysis of 1–4 linkages between *N*-acetyl- β -D-glucosamine and D-glucuronate residues in hyaluronate.

The long-term evolutionary conservation of such fundamental biochemical functions and their ubiquitous importance in the Kingdom Animalia provide the necessary functional basis for a body protein in one organism to perform a similar function when introduced into another. Consequently, the bioactivity of most toxins stems from one of the following three generalized mechanisms: (a) structural damage caused by catalyzing the hydrolysis of a universally present substrate, (b) physiological

imbalance or a short-term response caused by mimicking endogenous body proteins as if they were overexpressed, or (c) mimicking endogenous body proteins by acting competitive inhibitors to cause an opposite physiological imbalance or disruption of a physiological response. Hyaluronidase, PLA₂, and SMAse D represent examples of toxins that act according to the (a) mechanism because they catalyze the hydrolysis of universally present substrates. The binding of Bv8 or MIT-1 toxins on mammal prokineticin receptors and the hydrolytic release of kinins from kininogen by peptidase S1 enzymes (kallikrein subtype) are examples of the (b) mechanism. Finally, the inhibition of hemolymph host defense by Vn50, which acts as a competitive inhibitor of the peptidase S1 homologs from which it evolved, represents an example of the (c) mechanism. The effectiveness of the latter two mechanisms may be amplified if the toxin mimics the action of a pleiotropic protein that governs multiple downstream metabolic pathways or if the mimicked protein plays a central role in a complex protein cascade with feedback systems (e.g., thrombin-mimicking proteins affecting the blood coagulation process). Similar multifunctionality can be attained by mimicking multiple factors that share similar structures and biochemical activities. For example, peptidase S1 toxins in reptiles and *Blarina* short-tailed shrews mimic the different activities of the related kallikrein and thrombin, thereby lysing both kininogen (vasodilation) and fibrinogen (anticoagulation). As such, the toxicity of many venom proteins that interfere with hemostasis is enhanced by the fact that they are recruited from the same protein (super)families that constitute multiple components of the involved protein cascades. Such effects for each of these three scenarios described above allow the new venom-gland protein type to provide an immediately useful toxic bioactivity. Whereas the snake three-finger neurotoxins are derivations of α -neuropeptides similar to Lynx1/SLUR (54, 71), other neurotoxin types do not necessarily represent structural or even phylogenetic relatives of the ligands they mimic. However, many neurotoxins

act according to the second or third mechanism, either activating or deactivating ion channels or other receptors at neuromuscular junctions or in the central or peripheral nervous system.

Third, most toxins are recruited from body proteins involved in one or several short-term physiological processes. Because venoms are used for predation, competitor deterrence, or defense, they are effective only if they produce a rapid effect, such as fast immobilization of prey, instantaneous inhibition of blood coagulation in a host, or quick malaise or pain induction in a predator or competitor. For example, although the PK/PKR systems govern a large number of long-term physiological processes with potentially negative consequences for the host (e.g., predator anorexigenia caused by frog skin defensive AVIT peptides), the toxicity of AVIT peptides is largely determined by their short-term physiological effects (within minutes), which include powerful contraction of smooth muscle and hyperalgesia. Slow action on long-term physiological processes, lethal as they might be (e.g., accumulation of indigenous waste products or secondary systemic process after abnormal cellular depolarization), are unlikely to contribute much to the adaptive value of toxins because they are irrelevant to the basic ecological purposes of any given venom (although the anorexigenic effects have group level evolutionary benefits). This is not to say that some toxins do not have persistent effects, such as the long-term paralysis of parasitic wasps upon spider prey or, conversely, long-term paralysis by spiders of their prey. In these cases, the prey item is kept alive until the parasite eggs hatch or the spider eats its prey, respectively. Protein families that fulfill only strictly structural functions (such as keratins) or proteins that govern only time-consuming processes (such as cell growth or tissue differentiation factors) are therefore unlikely to be recruited as toxins. For a similar reason, toxin targeting is restricted to body parts that are readily accessible via the bloodstream, which also guarantees an acute effect. Most toxins additionally have an explicit extracellular substrate or site of

activity, even when they also affect intracellular processes.

Fourth, toxins seem to be recruited more frequently from body proteins with stable tertiary structures maintained by a high degree of disulfide cross-linking. The cysteine content of many animal toxins and their nontoxic relatives is remarkably high and evolutionarily conserved. Extensive disulfide cross-linking is a generalized feature of secretory proteins because it enhances molecular stability and protease resistance. In contrast, the correct folding of secreted globular enzymes is generally more sensitive to primary structure. A single mutation, such as the gain or loss of a proline or a crucial charge-state residue, may be enough to decimate the correct globular folding.

Fifth, once recruited as a functionally important component of the venom arsenal, the adaptive evolution of toxins is often reinforced by extensive gene duplication. Further duplication of the toxin gene is likely to be selectively favored because more gene copies could increase the toxin expression level in the gland and hence result in higher toxin doses and faster gland replenishment. The tandem duplication of gene copies provides the template, progressively increasing the chance for unequal crossing-over and gene conversion. This situation leads to the creation of a toxin multigene family in which differential mutations among paralogs allow them to obtain different levels of potency, functions, or complementary prey specificities (59, 81). The newly created toxin multigene families preserve the molecular scaffold of the ancestral protein but modify key surface functional residues to acquire a myriad of newly derived activities (38, 54, 59, 106). In globular enzymatic proteins, mutations are much more likely to interfere with correct post-translational folding than is the case with extensively cysteine cross-linked proteins. Thus, a negative selection pressure operates against the duplication and diversification of globular proteins, and they are instead overexpressed but with few new gene copies. For example, no new activities have been reported for toxins based

upon globular enzymes (e.g., hyaluronidase) as opposed to the extensive neofunctionalization of cysteine cross-linked enzymes (e.g., PLA₂).

One of the most intensively studied toxin structural classes, the disulfide-rich toxins of mass 3–5 kDa, which dominate the venoms of most spiders and venomous marine snails (144), appears to provide an exception to these general rules of toxin recruitment. The vast majority of these toxins contain the inhibitory cysteine knot (ICK) motif (also known as the knottin fold) that directs their three-dimensional fold (62, 120). Structurally similar peptides are also present in other arthropods and plants, where they function as either toxins targeting Ca²⁺ channels (scorpion and assassin bug toxins), antimicrobial agents (horseshoe crab tachystatin and plant thionins), or protease inhibitors (plant cyclotides). This highly compact domain (three disulfides within 20–40 amino acid residues) is quite scarce in vertebrates; the only known representative is the C-terminal domain of Agouti-related proteins, which are involved in feeding behavior and energy homeostasis (163). Based on the similarities in toxin prepropeptide transcript structure, it has been proposed that spider and cone snail ICK toxins are evolutionary related (176); however, their phylogenetic relationships, if any, are difficult to reconstruct with current methods because of their short and highly diverged sequences (36). Remote homology detection programs do not identify any spider toxin as a putative homolog if *Comus* sequences or multiple alignments are used as query, or vice versa. Some authors have proposed that the ICK is an elaboration of a simpler structural motif, the disulfide-directed β -hairpin (DDH), which is found in several vertebrate and invertebrate body proteins (159), and hence these toxins may have evolved from an ancestral DDH-containing body protein that was independently recruited into spider, scorpion, hymenopteran, and cone snail venoms. Nonetheless, convergent evolution of the three-dimensional fold—and its cysteine-pairing signature—cannot be ruled out because of the generalized scarcity of

information about nonvenom ICK-containing proteins in invertebrate genomes.

CONVERGENCE OF ACTION

In addition to convergence of the types of peptide/proteins recruited for use in the chemical arsenals, there has also been convergence of derived activities. Kunitz-type toxins represent a striking example of convergent toxin recruitment and convergent molecular evolution to produce the same derived activity. The derived venom forms in snakes (66), sea anemones (140), and cone snails (16) block K_V channels. Snake dendrotoxins fit the functional dyad motif found in several K^+ channel blockers (33), in which the minimum pharmacophore is defined by a critical basic residue (usually a lysine) located 7 Å from a hydrophobic residue. This functional site, assisted by different constellations of secondary residues for K^+ channel binding, is quite distinct from the protease inhibition site. Hence, the same fold is being used for two distinct functions, but different side chains are involved. Sea anemones contain a range of Kunitz-type protease inhibitors (10, 17, 69, 101), some of which are also dual action polypeptides (140). For example, Kaliclutidines 1 and -3 (also known as AsKC1-3) not only inhibit trypsin but also block K^+ channels. However, compared with the dendrotoxins, they are weaker inhibitors of $K_V1.2$, and compared with BPTI (bovine peptidic trypsin inhibitor), they are less potent trypsin inhibitors. *Conus striatus* venom contains a Kunitz-type neurotoxin that shares 33% amino acid sequence identity with BPTI and 35% with dendrotoxin I. This toxin, named Conkunitzin-S1, blocks *Shaker* potassium channels with an IC_{50} of approximately 60 nM. The Kunitz-type toxins in mygalomorph spider venoms also have a dual activity, acting as both trypsin inhibitors and K_V channel blockers (170). For example, huwentoxin-XI, a Kunitz-type toxin from the Chinese tarantula *Ornithoctonus huwena*, competitively inhibits trypsin with $K_i = 68$ nM (87), and it is also a weak blocker of $K_V1.1$ channels; the epitopes responsible for

protease inhibition and K_V channel block are spatially distinct and located at opposite ends of the cone-shaped molecule (170).

However, convergence of targeting also exists between derived forms of different toxin types. Notably, toxins from five distinct structural classes and phylogenetically distant organisms (CS $\alpha\beta$ short-chain scorpion toxins, Kunitz-type snake toxins, Kunitz-type and ICK snail toxins, and CRD-related sea anemone toxins) display the same kind of molecular determinants for K^+ channel recognition and blockage (34, 60).

Other neurological sites have also been convergently targeted at the neuromuscular junction (**Figure 7**). Virtually every site of action has been attacked on multiple occasions during evolution such as the neuronal nicotinic acetylcholine receptors convergently targeted by the κ -bungarotoxins (44) and also by conotoxins (85).

Targeting convergence can occur not only between lineages but also within a particular lineage. Two striking examples of targeting convergence within a particular lineage are the spider toxins on the one hand and the scorpion toxins on the other hand, which modulate sodium and potassium channels, respectively. The μ spider toxins are quite distinct from the δ spider toxins (such as the δ -atracotoxins, formerly known as robustoxin and versutoxin). The spider μ toxins, as in the case of the scorpion β toxins (40), cause a hyperpolarizing shift in the voltage-dependent activation of Na^+ channels, causing channel opening at lower potentials; consequently, they do not act as blockers or prolongers but more as channel activators (111). However, δ spider toxins delay Na^+ channel inactivation. Scorpion toxins that target K^+ channels act as current blockers. Most of them bind to the channel outer mouth with a homologous Lys residue plugging the ion-conduction pathway; however, some scorpion toxins do not obey this rule in spite of being bona fide blockers (38). Compelling evidence suggests that such toxins have evolved nonhomologous recognition sites that fulfill the same function (1) but through independent




Figure 7

Sites of neurotoxic convergence.

recruitment events from ancestral body forms with immune-related function (43).

Another example of neurotoxic targeting convergence within a lineage is snake toxins targeting the muscarinic receptors. The first of these to be described included three-finger toxins from *Dendroaspis* and *Naja* species (e.g., 22), but PLA₂ toxins with muscarinic actions were discovered later (70). Further, the *Dendroaspis* muscarinic three-finger toxins are not monophyletic (59), and thus represent a fascinating case of convergence within a single clade

of the same molecular scaffold being mutated two times for the same derived activity. *Dendroaspis* venoms are also extraordinary in convergently targeting L-type Ca²⁺ channels with derived forms of both Kunitz (169) and three-finger toxins (42).

Finally, sites of the coagulation cascade are also convergently targeted (Figure 8). Kunitz-domain proteins in snake venoms, for example, are similar to tick and biting midge hematophagous secretion proteins in mimicking the activity of endogenous TFPI (tissue




Figure 8

Sites of hematoxic convergence.

factor pathway inhibitor) by efficiently shutting down coagulation at the point of initiation. Using various protein scaffolds as templates, antiplatelet action mediated by the RGD tripeptide motif has also been a convergent target by snakes (74), leeches (83), and arthropods, including ticks (92) and horseflies (165). As with neurotoxicity, targeting convergence can occur within a single venomous clade, such as the RGD-containing distintegrins in viper venoms (74) and the RGD-containing three-finger peptides in mamba venoms (98).

WHAT DEFINES AN ANIMAL AS VENOMOUS OR A PROTEIN AS A TOXIN?

The convergent recruitment of different protein types for use as toxins raises the fundamental question of how to define an animal as venomous. Variables such as relative lethality to a prey item or to a potential predator are arbitrary and obscure evolutionary relationships. We reject the suggestion that rapid death should be the determinant of whether a toxic secretion should be classified as a venom. Apart from the difficulty in defining rapid death, such an overly restrictive and arbitrary definition obscures the evolutionary homology among the toxins, thus contributing little to our understanding of venom origin and evolution. The biological reality is complex, especially if we consider that the same secretion may have vastly different effects on different recipient animals. For example, the secretion of the Australian Paralysis Tick *Ixodes holocyclus* induces lethal paralytic neurotoxicity in humans as well as in a variety of non-native livestock and domestic animals, and thus clearly fits the conventional definition of venom. However, paralysis is not induced when these ticks feed on their natural hosts, including bandicoots, wallabies, and other marsupials (145). Tick hematophagous secretions contain a plethora of molecules that inhibit host hemostasis, immune response, and pain (so as to escape detection). A coevolutionary chemical arms race with native host animals has likely resulted in an especially potent secretion

in *I. holocyclus* to overcome resistant native prey. Therefore, non-native hosts are likely to be far more sensitive because they are immunologically naive. In addition, the susceptibility of individual hosts can be affected by prior history of exposure to ticks and seroconversion, so susceptible hosts can become resistant to paralysis (145). The tick, therefore, is confronted with a highly variable landscape of blood-meal hosts, ranging from highly susceptible to highly resistant. Because fitness is maximized when the tick can obtain a sufficient meal for reproduction regardless of the resistance of the host, selection favors the evolution of a potent pharmacopoeia, sufficient to allow feeding on resistant hosts but collaterally toxic to susceptible hosts.

Any venomous animal faces the same issues: Whether the venom is used for feeding or defense, it must work on the most resistant targets so effects will be more extreme on more susceptible animals. With this in mind, we regard venom as a secretion, produced in a specialized gland in one animal, and delivered to a target animal through the infliction of a wound (regardless of how tiny it) a venom must further contain molecules that disrupt normal physiological or biochemical processes so as to facilitate feeding or defense by the producing animal. By extension, toxins should be regarded as particular examples of intergenome active elements by means of their action on the extraorganismal space (Gene Ontology term number 0043245). This definition, based on biological function as opposed to an anthropocentric view of toxicity, recognizes that there is a vast range of effects of envenomation, from the hardly noticeable subversion of hemostatic defenses produced by a mosquito to the lethal effects of venomous snakes. Accordingly, we regard the feeding secretion of hematophagous specialists (e.g., arthropods or leeches) as a specialized subtype of venom.

Our broadened concept of venom is validated by the finding that many of the same classes of proteins are represented in the hematophagous secretions of blood-feeding arthropods and in more classical venoms of

reptiles and other organisms. Specifically, 11 of the 14 protein families discussed above are represented in at least one group of blood feeders, and nine of these families are found in two or more groups. In addition, protein families recruited in both conventional venom and hematophagous secretions of blood-feeding taxa often show the exact same biological activities. For example, a recently identified CRISP protein found in the buccal gland secretion of the sea lamprey (a blood parasite of fish) is not only homologous to CRISP toxins in toxicoferan reptiles, but also shows the same voltage-gated Ca^{2+} channel blocking activity and capacity to inhibit smooth muscle contraction (73). Despite the fact that this muscle contraction probably fulfills different adaptive roles in the two taxa (vasodilation to facilitate blood feeding in lamprey versus low blood pressure and prey immobilization in reptiles), they both meet the same overarching biological function: predation. Similar parallel recruitments in conventional venom systems and hematophagous sialomes are linked to the three major processes of hemostatic responses to vascular injury: vasoconstriction, platelet aggregation, and blood coagulation (**Figure 8**). Striking examples include (*a*) vasodilatory components (tachykinins, kallikreins, and natriuretic

peptides in various venomous animals versus sialokinins in mosquitoes), (*b*) inhibitors of ADP-induced platelet aggregation (apyrases in the venom of stinging insects and assassin bugs versus the hematophagous secretions of blood-feeding ticks, dipterans, fleas, and bugs), (*c*) platelet IIb–IIIa glycoprotein antagonists (snakes versus blood-feeding nematodes, ticks, and leeches), (*d*) thrombin inhibitors (viperid snakes versus leeches, ticks, dipterans, and triatomine bugs), (*e*) fibrino(geno)lytic proteases (toxicoferan reptiles and *Lonomia* caterpillars versus leeches), and (*f*) plasminogen activators (snakes versus vampire bats). Failure to recognize these numerous parallelisms between conventional venoms and secretions adapted to blood-feeding would be based on the arbitrary discrimination of their adaptive roles and would disregard their overruling similarities in terms of protein composition, bioactivity, and predatory function.

Acceptance of the broader definition proposed here expands our sample of venomous animals and increases the number of known independent occasions in which venom has evolved. This expansion of our sample size of venoms and venomous proteins will improve our understanding of factors underlying the evolution of venoms and their associated proteins.

SUMMARY POINTS

1. Convergetly recruited proteins share several conserved features: a secretory protein ancestor, functionally versatile protein ancestors with a fundamentally conserved basal activity, extensive disulfide cross-links, stable molecular scaffolds, and once recruited, adaptive evolution generates a suite of novel isoforms with neofunctionalization.
2. In addition to convergence of protein types utilized, convergence of toxic targeting has occurred extensively.
3. The convergence between venomous and hematophagous animals allows for an expanded definition of venom, thus recognizing the vast biodiversity of toxicity levels and types.

DISCLOSURE STATEMENT

The authors are not aware of any biases that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

This work was supported by grants to B.G.F. from the Australian Academy of Science, Australian French Association for Science and Technology, Australia and Pacific Science Foundation, CASS Foundation, Herman Slade Foundation, and University of Melbourne (Faculty of Medicine and Department of Biochemistry) and a Department of Innovation, Industry, and Regional Development Victoria Fellowship. The work was also supported by Australian Research Council Grants and an Australian Government Department of Education, Science, and Training/EGIDE International Science Linkages grant to B.G.F. and J.A.N.

LITERATURE CITED

1. Abdel-Mottaleb Y, Corzo G, Martin-Eauclaire MF, Satake H, Ceard B, et al. 2008. A common "hot spot" confers hERG blockade activity to α -scorpion toxins affecting K⁺ channels. *Biochem. Pharmacol.* 76:805–15
2. Alape-Giron A, Persson B, Cederlund E, Flores-Diaz M, Gutierrez JM, et al. 1999. Elapid venom toxins: multiple recruitments of ancient scaffolds. *Eur. J. Biochem.* 259:225–34
3. Amarant T, Burkhardt W, Levine H, Arochapaningo CL, Parikh I. 1991. Isolation and complete amino-acid-sequence of 2 fibrinolytic proteinases from the toxic saturnid caterpillar *Lonomia achelous*. *Biochim. Biophys. Acta* 1079:214–21
4. Ameri M, Wang X, Wilkerson MJ, Kanost MR, Broce AB. 2008. An immunoglobulin binding protein (Antigen 5) of the stable fly (Diptera: Muscidae) salivary gland stimulates bovine immune responses. *J. Med. Entomol.* 45:94–101
5. Andersen JF, Champagne DE, Weichsel A, Ribeiro JMC, Balfour CA, et al. 1997. Nitric oxide binding and crystallization of recombinant nitrophorin I, a nitric oxide transport protein from the blood-sucking bug *Rhodnius prolixus*. *Biochemistry* 36:4423–28
6. Andersen JF, Francischetti IMB, Valenzuela JG, Schuck P, Ribeiro JMC. 2003. Inhibition of hemostasis by a high affinity biogenic amine-binding protein from the saliva of a blood-feeding insect. *J. Biol. Chem.* 278:4611–17
7. Andersen JF, Gudderra NP, Francischetti IMB, Ribeiro JMC. 2005. The role of salivary lipocalins in blood feeding by *Rhodnius prolixus*. *Arch. Insect Biochem. Physiol.* 58:97–105
8. Andersen JF, Gudderra NP, Francischetti IMB, Valenzuela JG, Ribeiro JMC. 2004. Recognition of anionic phospholipid membranes by an antihemostatic protein from a blood-feeding insect. *Biochemistry* 43:6987–94
9. Andersen JF, Hinnebusch BJ, Lucas DA, Conrads TP, Veenstra TD, et al. 2007. An insight into the sialome of the oriental rat flea, *Xenopsylla cheopis* (Rots). *BMC Genomics* 8:102
10. Antuch W, Berndt KD, Chavez MA, Delfin J, Wuthrich K. 1993. The NMR solution structure of a kunitz-type proteinase-inhibitor from the sea-anemone *Stichodactyla helianthus*. *Eur. J. Biochem.* 212:675–84
11. Arca B, Lombardo F, Francischetti IMB, Pham VM, Mestres-Simon M, et al. 2007. An insight into the sialome of the adult female mosquito *Aedes albopictus*. *Insect Biochem. Mol. Biol.* 37:107–27
12. Arca B, Lombardo F, Valenzuela JG, Francischetti IMB, Marinotti O, et al. 2005. An updated catalogue of salivary gland transcripts in the adult female mosquito, *Anopheles gambiae*. *J. Exp. Biol.* 208:3971–86
13. Asgari S, Zhang GM, Zareie R, Schmidt O. 2003. A serine proteinase homolog venom protein from an endoparasitoid wasp inhibits melanization of the host hemolymph. *Insect Biochem. Mol. Biol.* 33:1017–24
14. Assumpcao TCF, Francischetti IMB, Andersen JF, Schwarz A, Santana JM, Ribeiro JMC. 2008. An insight into the sialome of the blood-sucking bug *Triatoma infestans*, a vector of Chagas' disease. *Insect Biochem. Mol. Biol.* 38:213–32
15. Batista CVF, Roman-Gonzalez SA, Salas-Castillo SP, Zamudio FZ, Gomez-Lagunas F, Possani LD. 2007. Proteomic analysis of the venom from the scorpion *Tityus stigmurus*: biochemical and physiological comparison with other Tityus species. *Comp. Biochem. Physiol. C-Toxicol. Pharmacol.* 146:147–57

16. Bayrhuber M, Vijayan V, Ferber M, Graf R, Korukottu J, et al. 2005. Conkunitzin-S1 is the first member of a new Kunitz-type neurotoxin family: structural and functional characterization. *J. Biol. Chem.* 280:23766–70
17. Beress L. 1982. Biologically-active compounds from coelenterates. *Pure Appl. Chem.* 54:1981–94
18. Binford GJ, Cordes MHJ, Wells MA. 2005. Sphingomyelinase D from venoms of *Loxosceles* spiders: evolutionary insights from cDNA sequences and gene structure. *Toxicon* 45:547–60
19. Binford GJ, Wells MA. 2003. The phylogenetic distribution of sphingomyelinase D activity in venoms of *Haplogygne* spiders. *Comp. Biochem. Physiol. B-Biochem. Mol. Biol.* 135:25–33
20. Birkemo GA, Luders T, Andersen O, Nes IF, Nissen-Meyer J. 2003. Hipposin, a histone-derived antimicrobial peptide in Atlantic halibut (*Hippoglossus hippoglossus* L.). *Biochim. Biophys. Acta-Proteins Proteomics* 1646:207–15
21. Boisbouvier J, Albrand JP, Blackledge M, Jaquinod M, Schweitz H, et al. 1998. A structural homologue of colipase in black mamba venom revealed by NMR floating disulphide bridge analysis. *J. Mol. Biol.* 283:205–19
22. Bradley KN. 2000. Muscarinic toxins from the green mamba. *Pharmacol. Ther.* 85:87–109
23. Brown RL, Haley TL, West KA, Crabb JW. 1999. Pseudechetoxin: a peptide blocker of cyclic nucleotide-gated ion channels. *Proc. Natl. Acad. Sci. USA* 96:754–59
24. Brown RL, Lynch LL, Haley TL, Arsanjani R. 2003. Pseudechetoxin binds to the pore turret of cyclic nucleotide-gated ion channels. *J. Gen. Physiol.* 122:749–60
25. Campbell CL, Vandyke KA, Letchworth GJ, Drolet BS, Hanekamp T, Wilson WC. 2005. Midgut and salivary gland transcriptomes of the arbovirus vector *Culicoides sonorensis* (Diptera: Ceratopogonidae). *Insect Mol. Biol.* 14:121–36
26. Campbell CL, Wilson WC, Manninen K. 2005. Characterization of differentially expressed midgut genes in orbivirus vector populations. *Am. J. Trop. Med. Hyg.* 73:144
27. Champagne DE, Nussenzweig RH, Ribeiro JMC. 1995. Purification, partial characterization, and cloning of nitric oxide-carrying heme-proteins (nitrophorins) from salivary glands of the blood-sucking insect *Rhodnius prolixus*. *J. Biol. Chem.* 270:8691–95
28. Charlab R, Valenzuela JG, Rowton ED, Ribeiro JMC. 1999. Toward an understanding of the biochemical and pharmacological complexity of the saliva of a hematophagous sand fly *Lutzomyia longipalpis*. *Proc. Natl. Acad. Sci. USA* 96:15155–60
29. Chen DS, Kini RM, Yuen R, Khoo HE. 1997. Haemolytic activity of stonustoxin from stonefish (*Synanceja horrida*) venom: pore formation and the role of cationic amino acid residues. *Biochem. J.* 325:685–91
30. Chen TB, Farragher S, Bjourson AJ, Orr DF, Rao P, Shaw C. 2003. Granular gland transcriptomes in stimulated amphibian skin secretions. *Biochem. J.* 371:125–30
31. Chen TB, Xue YZ, Zhou M, Shaw C. 2005. Molecular cloning of mRNA from toad granular gland secretion and lyophilized skin: identification of Bo8—a novel prokineticin from *Bombina orientalis*. *Peptides* 26:377–83
32. Cordes MHJ, Binford GJ. 2006. Lateral gene transfer of a dermonecrotic toxin between spiders and bacteria. *Bioinformatics* 22:264–68
33. Dauplais M, Gilquin B, Possani LD, GurrolaBriones G, Roumestand C, Menez A. 1995. Determination of the three-dimensional solution structure of noxiustoxin: analysis of structural differences with related short-chain scorpion toxins. *Biochemistry* 34:16563–73
34. Dauplais M, Lecoq A, Song JX, Cotton J, Jamin N, et al. 1997. On the convergent evolution of animal toxins: conservation of a diad of functional residues in potassium channel-blocking toxins with unrelated structures. *J. Biol. Chem.* 272:4302–9
35. Davidson FF, Dennis EA. 1990. Evolutionary relationships and implications for the regulation of phospholipase A₂ from snake venom to human secreted forms. *J. Mol. Evol.* 31:228–38
36. De La Vega RCR. 2005. A note on the evolution of spider toxins containing the ICK-motif. *Toxin Rev.* 24:385–97
37. de la Vega RCR, Garcia BI, D'Ambrosio C, Diego-Garcia E, Scaloni A, Possani LD. 2004. Antimicrobial peptide induction in the haemolymph of the Mexican scorpion *Centruroides limpidus limpidus* in response to septic injury. *Cell. Mol. Life Sci.* 61:1507–19

38. de la Vega RCR, Merino E, Becerril B, Possani LD. 2003. Novel interactions between K⁺ channels and scorpion toxins. *Trends Pharmacol. Sci.* 24:222–27
39. de la Vega RCR, Possani LD. 2005. On the evolution of invertebrate defensins. *Trends Genet.* 21:330–32
40. de la Vega RCR, Possani LD. 2007. Novel paradigms on scorpion toxins that affects the activating mechanism of sodium channels. *Toxicon* 49:171–80
41. de Plater GM, Martin RL, Milburn PJ. 1998. A C-type natriuretic peptide from the venom of the platypus (*Ornithorhynchus anatinus*): structure and pharmacology. *Comp. Biochem. Physiol. C-Toxicol. Pharmacol.* 120:99–110
42. de Weille JR, Schweitz H, Maes P, Tartar A, Lazdunski M. 1991. Calciseptine, a peptide isolated from black mamba venom, is a specific blocker of the L-type calcium channel. *Proc. Natl. Acad. Sci. USA* 88:2437–40
43. Diego-Garcia E, Abdel-Mottaleb Y, Schwartz EF, de la Vega RCR, Tytgat J, Possani LD. 2008. Cytolytic and K⁺ channel blocking activities of β -KTx and scorpine-like peptides purified from scorpion venoms. *Cell. Mol. Life Sci.* 65:187–200
44. Dryer SE, Chiappinelli VA. 1983. Kappa-bungarotoxin: an intracellular study demonstrating blockade of neuronal nicotinic receptors by a snake neurotoxin. *Brain Res.* 289:317–21
45. Dubin G. 2005. Proteinaceous cysteine protease inhibitors. *Cell. Mol. Life Sci.* 62:653–69
46. Ehret-Sabatier L, Loew D, Goyffon M, Fehlbaum P, Hoffmann JA, et al. 1996. Characterization of novel cysteine-rich antimicrobial peptides from scorpion blood. *J. Biol. Chem.* 271:29537–44
47. Fang KSY, Vitale M, Fehlner P, King TP. 1988. cDNA cloning and primary structure of a white-face hornet venom allergen, antigen 5. *Proc. Natl. Acad. Sci. USA* 85:895–99
48. Filippovich I, Sorokina N, Masci PP, de Jersey J, Whitaker AN, et al. 2002. A family of textilinin genes, two of which encode proteins with antihemorrhagic properties. *Br. J. Haematol.* 119:376–84
49. Fox JW, Serrano SMT. 2008. Exploring snake venom proteomes: multifaceted analyses for complex toxin mixtures. *Proteomics* 8:909–20
50. Francischetti IMB, Andersen JF, Ribeiro JMC. 2002. Biochemical and functional characterization of recombinant *Rhodnius prolixus* platelet aggregation inhibitor 1 as a novel lipocalin with high affinity for adenosine diphosphate and other adenine nucleotides. *Biochemistry* 41:3810–18
51. Francischetti IMB, Ribeiro JMC, Champagne D, Andersen J. 2000. Purification, cloning, expression, and mechanism of action of a novel platelet aggregation inhibitor from the salivary gland of the blood-sucking bug, *Rhodnius prolixus*. *J. Biol. Chem.* 275:12639–50
52. Francischetti IMB, Valenzuela JG, Andersen JF, Mather TN, Ribeiro JMC. 2002. Ixolaris, a novel recombinant tissue factor pathway inhibitor (TFPI) from the salivary gland of the tick, *Ixodes scapularis*: identification of factor X and factor Xa as scaffolds for the inhibition of factor VIIa/tissue factor complex. *Blood* 99:3602–12
53. Froy O, Gurevitz M. 1998. Membrane potential modulators: a thread of scarlet from plants to humans. *EASEB J.* 12:1793–96
54. Fry BG. 2005. From genome to “venome”: molecular origin and evolution of the snake venom proteome inferred from phylogenetic analysis of toxin sequences and related body proteins. *Genome Res.* 15:403–20
55. Fry BG, Scheib H, Van Der Weerd L, Young B, McNaughtan J, et al. 2008. Evolution of an arsenal. *Mol. Cell. Proteomics* 7:215–46
56. Fry BG, Vidal N, Norman JA, Vonk FJ, Scheib H, et al. 2006. Early evolution of the venom system in lizards and snakes. *Nature* 439:584–88
57. Fry BG, Wickramaratana JC, Lemme S, Beuve A, Garbers D, et al. 2005. Novel natriuretic peptides from the venom of the inland taipan (*Oxyuranus microlepidotus*): isolation, chemical and biological characterisation. *Biochem. Biophys. Res. Commun.* 327:1011–15
58. Fry BG, Wuster W. 2004. Assembling an arsenal: origin and evolution of the snake venom proteome inferred from phylogenetic analysis of toxin sequences. *Mol. Biol. Evol.* 21:870–83
59. Fry BG, Wuster W, Kini RM, Brusic V, Khan A, et al. 2003. Molecular evolution and phylogeny of elapid snake venom three-finger toxins. *J. Mol. Evol.* 57:110–29
60. Gasparini S, Gilquin B, Menez A. 2004. Comparison of sea anemone and scorpion toxins binding to Kv1 channels: an example of convergent evolution. *Toxicon* 43:901–8

61. Gmachl M, Kreil G. 1993. Bee venom hyaluronidase is homologous to a membrane-protein of mammalian sperm. *Proc. Natl. Acad. Sci. USA* 90:3569–73
62. Gracy J, Le-Nguyen D, Gelly JC, Kaas Q, Heitz A, Chiche L. 2008. KNOTTIN: the knottin or inhibitor cystine knot scaffold in 2007. *Nucleic Acids Res.* 36:D314–19
63. Grunclova L, Horn M, Vancova M, Sojka D, Franta Z, et al. 2006. Two secreted cystatins of the soft tick *Ornithodoros moubata*: differential expression pattern and inhibitory specificity. *Biol. Chem.* 387:1635–44
64. Gudderra NP, Ribeiro JMC, Andersen JF. 2005. Structural determinants of factor IX(a) binding in nitrophorin 2, a lipocalin inhibitor of the intrinsic coagulation pathway. *J. Biol. Chem.* 280:25022–28
65. Harrison RA, Ibson F, Wilbraham D, Wagstaff SC. 2007. Identification of cDNAs encoding viper venom hyaluronidases: cross-generic sequence conservation of full-length and unusually short variant transcripts. *Gene* 392:22–33
66. Harvey AL, Robertson B. 2004. Dendrotoxins: structure-activity relationships and effects on potassium ion channels. *Curr. Med. Chem.* 11:3065–72
67. Hawdon JM, Jones BF, Hoffman DR, Hotez PJ. 1996. Cloning and characterization of *Ancylostoma* secreted protein: a novel protein associated with the transition to parasitism by infective hookworm larvae. *J. Biol. Chem.* 271:6672–78
68. Hisada M, Satake H, Masuda K, Aoyama M, Murata K, et al. 2005. Molecular components and toxicity of the venom of the solitary wasp, *Anoplius samariensis*. *Biochem. Biophys. Res. Commun.* 330:1048–54
69. Honma T, Shiomi K. 2006. Peptide toxins in sea anemones: structural and functional aspects. *Mar. Biotechnol.* 8:1–10
70. Huang LF, Zheng HB, Xu Y, Song HT, Yu CX. 2008. A snake venom phospholipase A(2) with high affinity for muscarinic acetylcholine receptors acts on guinea pig ileum. *Toxicon* 51:1008–16
71. Ibanez-Tallon I, Miwa JM, Wang HL, Adams NC, Crabtree GW, et al. 2002. Novel modulation of neuronal nicotinic acetylcholine receptors by association with the endogenous prototoxin lynx1. *Neuron* 33:893–903
72. Inceoglu B, Lango J, Jing J, Chen LL, Doymaz F, et al. 2003. One scorpion, two venoms: pre venom of *Parabuthus transvaalicus* acts as an alternative type of venom with distinct mechanism of action. *Proc. Natl. Acad. Sci. USA* 100:922–27
73. Ito N, Mita M, Takahashi Y, Matsushima A, Watanabe YG, et al. 2007. Novel cysteine-rich secretory protein in the buccal gland secretion of the parasitic lamprey, *Lethenteron japonicum*. *Biochem. Biophys. Res. Commun.* 358:35–40
74. Kamiguti AS, Zuzel M, Theakston RDG. 1998. Snake venom metalloproteinases and disintegrins: interactions with cells. *Braz. J. Med. Biol. Res.* 31:853–62
75. Kato H, Anderson JM, Kamhawi S, Oliveira F, Lawyer PG, et al. 2006. High degree of conservancy among secreted salivary gland proteins from two geographically distant *Plebotomus duboscqi* sandflies populations (Mali and Kenya). *BMC Genomics* 7:226
76. Kim HS, Park CB, Kim MS, Kim SC. 1996. cDNA cloning and characterization of buforin I, an antimicrobial peptide: a cleavage product of histone H2A. *Biochem. Biophys. Res. Commun.* 229:381–87
77. King TP, Lu G, Gonzalez M, Qian NF, Soldatova L. 1996. Yellow jacket venom allergens, hyaluronidase and phospholipase: sequence similarity and antigenic cross-reactivity with their hornet and wasp homologs and possible implications for clinical allergy. *J. Allergy Clin. Immunol.* 98:588–600
78. Kini RM. 2003. Excitement ahead: structure, function and mechanism of snake venom phospholipase A₂ enzymes. *Toxicon* 42:827–40
79. Kita M, Nakamura Y, Okumura Y, Ohdachi SD, Oba Y, et al. 2004. Blarina toxin, a mammalian lethal venom from the short-tailed shrew *Blarina brevicauda*: isolation and characterization. *Proc. Natl. Acad. Sci. USA* 101:7542–47
80. Kita M, Okumura Y, Ohdachi SD, Oba Y, Yoshikuni M, et al. 2005. Purification and characterisation of blarinasin, a new tissue kallikrein-like protease from the short-tailed shrew *Blarina brevicauda*: comparative studies with blarina toxin. *Biol. Chem.* 386:177–82
81. Kordis D, Gubenek F. 2000. Adaptive evolution of animal toxin multigene families. *Gene* 261:43–52
82. Kratzschmar J, Haendler B, Langer G, Boidol W, Bringmann P, et al. 1991. The plasminogen activator family from the salivary-gland of the vampire bat *Desmodus rotundus*: cloning and expression. *Gene* 105:229–37

83. Krezel AM, Ulmer JS, Wagner G, Lazarus RA. 2000. Recombinant decorsin: dynamics of the RGD recognition site. *Protein Sci.* 9:1428–38
84. Krishnan A, Nair PN, Jones D. 1994. Isolation, cloning, and characterization of new chitinase stored in active form in chitin-lined venom reservoir. *J. Biol. Chem.* 269:20971–76
85. Lewis RJ. 2004. Conotoxins as selective inhibitors of neuronal ion channels, receptors and transporters. *IUBMB Life* 56:89–93
86. Li S, Kwon J, Aksoy S. 2001. Characterization of genes expressed in the salivary glands of the tsetse fly, *Glossina morsitans morsitans*. *Insect Mol. Biol.* 10:69–76
87. Liang SP. 2004. An overview of peptide toxins from the venom of the Chinese bird spider *Selenocosmia buwena* Wang [= *Ornithoctonus buwena* (Wang)]. *Toxicon* 43:575–85
88. Low KSY, Gwee MCE, Yuen R, Gopalakrishnakone P, Khoo HE. 1994. Stonustoxin: effects on neuromuscular function in vitro and in vivo. *Toxicon* 32:573–81
89. Magalhaes GS, Junqueira-de-Azevedo ILM, Lopes-Ferreira M, Lorenzini DM, Ho PL, Moura-da-Silva AM. 2006. Transcriptome analysis of expressed sequence tags from the venom glands of the fish *Thalassophryne nattereri*. *Biochimie* 88:693–99
90. Mans BJ, Andersen JF, Francischetti IMB, Valenzuel JG, Schwan TG, et al. 2008. Comparative sialomics between hard and soft ticks: implications for the evolution of blood-feeding behavior. *Insect Biochem. Mol. Biol.* 38:42–58
91. Mans BJ, Andersen JF, Schwan TG, Ribeiro JMC. 2008. Characterization of anti-hemostatic factors in the argasid, *Argas monolakensis*: implications for the evolution of blood-feeding in the soft tick family. *Insect Biochem. Mol. Biol.* 38:22–41
92. Mans BJ, Louw AI, Neitz AWH. 2002. Savignygrin, a platelet aggregation inhibitor from the soft tick *Ornithodoros savignyi*, presents the RGD integrin recognition motif on the Kunitz-BPTI fold. *J. Biol. Chem.* 277:21371–78
93. Mans BJ, Louw AI, Neitz AWH. 2003. The major tick salivary gland proteins and toxins from the soft tick, *Ornithodoros savignyi*, are part of the tick lipocalin family: implications for the origins of tick toxicoses. *Mol. Biol. Evol.* 20:1158–67
94. Deleted in proof
95. Mans BJ, Ribeiro JMC. 2008. A novel clade of cysteinyl leukotriene scavengers in soft ticks. *Insect Biochem. Mol. Biol.* 38:862–70
96. Mans BJ, Ribeiro JMC. 2008. Function, mechanism and evolution of the moubatin-clade of soft tick lipocalins. *Insect Biochem. Mol. Biol.* 38:841–52
97. Mans BJ, Ribeiro JMC, Andersen JF. 2008. Structure, function, and evolution of biogenic amine-binding proteins in soft ticks. *J. Biol. Chem.* 283:18721–33
98. McDowell RS, Dennis MS, Louie A, Shuster M, Mulkerrin MG, Lazarus RA. 1992. Mambin, a potent glycoprotein-IIb-IIIa antagonist and platelet-aggregation inhibitor structurally related to the short neurotoxins. *Biochemistry* 31:4766–72
99. Menendez A, Finlay BB. 2007. Defensins in the immunology of bacterial infections. *Curr. Opin. Immunol.* 19:385–91
100. Milne TJ, Abbenante G, Tyndall JDA, Halliday J, Lewis RJ. 2003. Isolation and characterization of a cone snail protease with homology to CRISP proteins of the pathogenesis-related protein superfamily. *J. Biol. Chem.* 278:31105–10
101. Minagawa S, Ishida M, Shimakura K, Nagashima Y, Shiomi K. 1997. Isolation and amino acid sequences of two Kunitz-type protease inhibitors from the sea anemone *Anthopleura* aff. *xanthogrammica*. *Comp. Biochem. Physiol. B-Biochem. Mol. Biol.* 118:381–86
102. Mollay C, Wechselberger C, Mignogna G, Negri L, Melchiorri P, et al. 1999. Bv8, a small protein from frog skin and its homologue from snake venom induce hyperalgesia in rats. *Eur. J. Pharmacol.* 374:189–96
103. Morita A, Isawa H, Orito Y, Iwanaga S, Chinzei Y, Yuda M. 2006. Identification and characterization of a collagen-induced platelet aggregation inhibitor, triplatin, from salivary glands of the assassin bug, *Triatoma infestans*. *FEBS J.* 273:2955–62
104. Morita T. 2005. Structures and functions of snake venom CLPs (C-type lectin-like proteins) with anticoagulant-, procoagulant-, and platelet-modulating activities. *Toxicon* 45:1099–114

105. Morrissette J, Kratzschmar J, Haendler B, Elhayek R, Mochcamorales J, et al. 1995. Primary structure and properties of helothermine, a peptide toxin that blocks ryanodine receptors. *Biophys. J.* 68:2280–88
106. Mouhat S, Jouirou B, Mosbah A, De Waard M, Sabatier JM. 2004. Diversity of folds in animal toxins acting on ion channels. *Biochem. J.* 378:717–26
107. Nagaraju S, Devaraja S, Kemparaju K. 2007. Purification and properties of hyaluronidase from *Hippasa partita* (funnel web spider) venom gland extract. *Toxicon* 50:383–93
108. Negri L, Lattanzi R, Giannini E, Melchiorri P. 2007. Bv8/prokineticin proteins and their receptors. *Life Sci.* 81:1103–16
109. Nevalainen TJ. 2008. Phospholipases A₂ in the genome of the sea anemone *Nematostella vectensis*. *Comp. Biochem. Physiol. D-Genomics Proteomics* 3:226–33
110. Ng HC, Ranganathan S, Chua KL, Khoo HE. 2005. Cloning and molecular characterization of the first aquatic hyaluronidase, SFHYA1, from the venom of stonefish (*Synanceja horrida*). *Gene* 346:71–81
111. Nicholson GM. 2007. Insect-selective spider toxins targeting voltage-gated sodium-channels. *Toxicon* 49:490–512
112. Nobile M, Magnelli V, Lagostena L, Mochcamorales J, Possani LD, Prestipino G. 1994. The toxin helothermine affects potassium currents in newborn rat cerebellar granule cells. *J. Membr. Biol.* 139:49–55
113. Nobile M, Noceti F, Prestipino G, Possani LD. 1996. Helothermine, a lizard venom toxin, inhibits calcium current in cerebellar granules. *Exp. Brain Res.* 110:15–20
114. Noeske-Jungblut C, Haendler B, Donner P, Alagon A, Possani L, Schleuning WD. 1995. Triabin, a highly potent exosite inhibitor of thrombin. *J. Biol. Chem.* 270:28629–34
115. Noeske-Jungblut C, Kratzschmar J, Haendler B, Alagon A, Possani L, et al. 1994. An inhibitor of collagen-induced platelet-aggregation from the saliva of *Triatoma pallidipennis*. *J. Biol. Chem.* 269:5050–53
116. Nunn MA, Sharma A, Paesen GC, Adamson S, Lissina O, et al. 2005. Complement inhibitor of C5 activation from the soft tick *Ornithodoros moubata*. *J. Immunol.* 174:2084–91
117. Olivera BM, Teichert RW. 2007. Diversity of the neurotoxic *Conus* peptides: a model for concerted pharmacological discovery. *Mol. Interv.* 7:251–60
118. Paesen GA, Adams PL, Nuttall PA, Stuart DL. 2000. Tick histamine-binding proteins: lipocalins with a second binding cavity. *Biochim. Biophys. Acta-Protein Struct. Mol. Enzymol.* 1482:92–101
119. Paesen GC, Adams PL, Harlos K, Nuttall PA, Stuart DI. 1999. Tick histamine-binding proteins: isolation, cloning, and three-dimensional structure. *Mol. Cell* 3:661–71
120. Pallyghy PK, Nielsen KJ, Craik DJ, Norton RS. 1994. A common structural motif incorporating a cystine knot and a triple-stranded β -sheet in toxic and inhibitory polypeptides. *Protein Sci.* 3:1833–39
121. Park IY, Park CB, Kim MS, Kim SC. 1998. Parasin I, an antimicrobial peptide derived from histone H2A in the catfish, *Parasilurus asotus*. *FEBS Lett.* 437:258–62
122. Patthy L. 2003. Modular assembly of genes and the evolution of new functions. *Genetica* 118:217–31
123. Possani LD, Becerril B, Delepierre M, Tytgat J. 1999. Scorpion toxins specific for Na⁺-channels. *Eur. J. Biochem.* 264:287–300
124. Possani LD, Merino E, Corona M, Bolivar F, Becerril B. 2000. Peptides and genes coding for scorpion toxins that affect ion-channels. *Biochimie* 82:861–68
125. Pung YF, Wong PTH, Kumar PP, Hodgson WC, Kini RM. 2005. Ohanin, a novel protein from king cobra venom, induces hypolocomotion and hyperalgesia in mice. *J. Biol. Chem.* 280:13137–47
126. Pungercar J, Krizaj I. 2007. Understanding the molecular mechanism underlying the presynaptic toxicity of secreted phospholipases A₂. *Toxicon* 50:871–92
127. Reis CV, Andrade SA, Ramos OHP, Ramos CRR, Ho PL, et al. 2006. Lopap, a prothrombin activator from *Lonomia obliqua* belonging to the lipocalin family: recombinant production, biochemical characterization and structure-function insights. *Biochem. J.* 398:295–302
128. Ribeiro JMC, Alarcon-Chaidez F, Francischetti IMB, Mans BJ, Mather TN, et al. 2006. An annotated catalog of salivary gland transcripts from *Ixodes scapularis* ticks. *Insect Biochem. Mol. Biol.* 36:111–29
129. Ribeiro JMC, Andersen J, Silva-Neto MAC, Pham VM, Garfield MK, Valenzuela JG. 2004. Exploring the sialome of the blood-sucking bug *Rhodnius prolixus*. *Insect Biochem. Mol. Biol.* 34:61–79

130. Ribeiro JMC, Arca B, Lombardo F, Calvo E, Phan VM, et al. 2007. An annotated catalogue of salivary gland transcripts in the adult female mosquito, *Aedes aegypti*. *BMC Genomics* 8:6
131. Ribeiro JMC, Charlab R, Pham VM, Garfield M, Valenzuela JG. 2004. An insight into the salivary transcriptome and proteome of the adult female mosquito *Culex pipiens quinquefasciatus*. *Insect Biochem. Mol. Biol.* 34:543–63
132. Ribeiro JMC, Schneider M, Guimaraes JA. 1995. Purification and characterization of prolixin-S (nitrophorin 2), the salivary anticoagulant of the bloodsucking bug *Rhodnius prolixus*. *Biochem. J.* 308:243–49
133. Ribeiro JMC, Walker FA. 1994. High-affinity histamine-binding and antihistaminic activity of the salivary nitric oxide-carrying heme protein (nitrophorin) of *Rhodnius prolixus*. *J. Exp. Med.* 180:2251–57
134. Ricci-Silva ME, Valente RH, Leon IR, Tambourgi DV, Ramos OHP, et al. 2008. Immunochemical and proteomic technologies as tools for unravelling toxins involved in envenoming by accidental contact with *Lonomia obliqua* caterpillars. *Toxicon* 51:1017–28
135. Sangamnatdej S, Paesen GC, Slovak M, Nuttall PA. 2002. A high affinity serotonin- and histamine-binding lipocalin from tick saliva. *Insect Mol. Biol.* 11:79–86
136. Santos A, Ribeiro JMC, Lehane MJ, Gontijo NF, Veloso AB, et al. 2007. The sialotranscriptome of the blood-sucking bug *Triatoma brasiliensis* (Hemiptera, Triatominae). *Insect Biochem. Mol. Biol.* 37:702–12
137. Schaloske RH, Dennis EA. 2006. The phospholipase A₂ superfamily and its group numbering system. *Biochim. Biophys. Acta-Mol. Cell Biol. Lipids* 1761:1246–59
138. Schleuning WD, Alagon A, Boidol W, Bringmann P, Petri T, et al. 1992. Plasminogen activators from the saliva of *Desmodus rotundus* (common vampire bat): unique fibrin specificity. *Ann. N. Y. Acad. Sci.* 667:395–403
139. Schwartz EF, Diego-Garcia E, de la Vega RCR, Possani LD. 2007. Transcriptome analysis of the venom gland of the Mexican scorpion *Hadrurus gertschi* (Arachnida: Scorpiones). *BMC Genomics* 8:116
140. Schweitz H, Bruhn T, Guillemare E, Moinier D, Lancelin JM, et al. 1995. Kalicludines and kaliseptine: Two different classes of sea-anemone toxins for voltage-sensitive K⁺ channels. *J. Biol. Chem.* 270:25121–26
141. Schweitz H, Heurteaux C, Bois P, Moinier D, Romey G, Lazdunski M. 1994. Calcicludine, a venom peptide of the kunitz-type protease inhibitor family, is a potent blocker of high-threshold Ca²⁺ channels with a high-affinity for L-type channels in cerebellar granule neurons. *Proc. Natl. Acad. Sci. USA* 91:878–82
142. Schweitz H, Pacaud P, Diochot S, Moinier D, Lazdunski M. 1999. MIT1, a black mamba toxin with a new and highly potent activity on intestinal contraction. *FEBS Lett.* 461:183–88
143. Soares MR, Oliveira-Carvalho AL, Wermelinger LS, Zingali RB, Ho PL, et al. 2005. Identification of novel bradykinin-potentiating peptides and C-type natriuretic peptide from *Lachesis muta* venom. *Toxicon* 46:31–38
144. Sollod BL, Wilson D, Zhaxybayeva O, Gogarten JP, Drinkwater R, King GF. 2005. Were arachnids the first to use combinatorial peptide libraries? *Peptides* 26:131–39
145. Stone BF. 1988. Tick paralysis, particularly involving *Ixodes holocyclus* and other *Ixodes* species. In *Advances in Disease Vector Research*, ed. KF Harris, 5:61–85. New York: Springer-Verlag
146. Sung JML, Low KSY, Khoo HE. 2002. Characterization of the mechanism underlying stonustoxin-mediated relaxant response in the rat aorta in vitro. *Biochem. Pharmacol.* 63:1113–18
147. Sutherland S, Lane WR. 1969. Toxins and mode of envenomation of common ringed or blue-banded octopus. *Med. J. Aust.* 1:893–97
148. Szeto TH, Wang XH, Smith R, Connor M, Christie MJ, et al. 2000. Isolation of a funnel-web spider polypeptide with homology to mamba intestinal toxin 1 and the embryonic head inducer Dickkopf-1. *Toxicon* 38:429–42
149. Tamiya T, Fujimi TJ. 2006. Molecular evolution of toxin genes in Elapidae snakes. *Mol. Divers.* 10:529–43
150. Todd SM, Sonenshine DE, Hynes WL. 2007. Tissue and life-stage distribution of a defensin gene in the Lone Star tick, *Amblyomma americanum*. *Med. Vet. Entomol.* 21:141–47
151. Torres AM, Kuchel PW. 2004. The β-defensin-fold family of polypeptides. *Toxicon* 44:581–88
152. Torres AM, Wang XH, Fletcher JI, Alewood D, Alewood PF, et al. 1999. Solution structure of a defensin-like peptide from platypus venom. *Biochem. J.* 341:785–94

153. Tu AT, Hendon RR. 1983. Characterization of lizard venom hyaluronidase and evidence for its action as a spreading factor. *Comp. Biochem. Physiol. B-Biochem. Mol. Biol.* 76:377–83
154. Valenzuela JG, Garfield M, Rowton ED, Pham VM. 2004. Identification of the most abundant secreted proteins from the salivary glands of the sand fly *Lutzomyia longipalpis*, vector of *Leishmania chagasi*. *J. Exp. Biol.* 207:3717–29
155. Veiga ABG, Ribeiro JMC, Guimaraes JA, Francischetti IMB. 2005. A catalog for the transcripts from the venomous structures of the caterpillar *Lonomia obliqua*: identification of the proteins potentially involved in the coagulation disorder and hemorrhagic syndrome. *Gene* 355:11–27
156. Vonk FJ, Admiraal JF, Jackson K, Reshef R, de Bakker MAG, et al. 2008. Evolutionary origin and development of snake fangs. *Nature* 454:630–33
157. Wagstaff SC, Favreau P, Cheneval O, Laing GD, Wilkinson MC, et al. 2008. Molecular characterisation of endogenous snake venom metalloproteinase inhibitors. *Biochem. Biophys. Res. Commun.* 365:650–56
158. Wang J, Shen B, Guo M, Lou XH, Duan YY, et al. 2005. Blocking effect and crystal structure of natriin toxin, a cysteine-rich secretory protein from *Naja atra* venom that targets the BK_{Ca} channel. *Biochemistry* 44:10145–52
159. Wang XH, Connor M, Smith R, Maciejewski MW, Howden MEH, et al. 2000. Discovery and characterization of a family of insecticidal neurotoxins with a rare vicinal disulfide bridge. *Nat. Struct. Biol.* 7:505–13
160. Waxman L, Connolly TM. 1993. Isolation of an inhibitor selective for collagen-stimulated platelet-aggregation from the soft tick *Ornithodoros moubata*. *J. Biol. Chem.* 268:5445–49
161. Wei AZ, Alexander RS, Duke J, Ross H, Rosenfeld SA, Chang CH. 1998. Unexpected binding mode of tick anticoagulant peptide complexed to bovine factor Xa. *J. Mol. Biol.* 283:147–54
162. Wen SP, Wilson DTR, Kuruppu S, Korsinczky MLJ, Hedrick J, et al. 2005. Discovery of an MIT-like atracotoxin family: spider venom peptides that share sequence homology but not pharmacological properties with AVIT family proteins. *Peptides* 26:2412–26
163. Wilczynski AM, Joseph CG, Haskell-Luevano C. 2005. Current trends in the structure activity relationship studies of the endogenous agouti-related protein (AGRP) melanocortin receptor antagonist. *Med. Res. Rev.* 25:545–56
164. Wong JH, Xia LX, Ng TB. 2007. A review of defensins of diverse origins. *Curr. Protein Pept. Sci.* 8:446–59
165. Xu XQ, Yang HL, Ma DY, Wu J, Wang YP, et al. 2008. Toward an understanding of the molecular mechanism for successful blood feeding by coupling proteomics analysis with pharmacological testing of horsefly salivary glands. *Mol. Cell. Proteomics* 7:582–90
166. Xu Y, Bruno JF, Luft BJ. 2005. Identification of novel tick salivary gland proteins for vaccine development. *Biochem. Biophys. Res. Commun.* 326:901–4
167. Yamazaki Y, Brown RL, Morita T. 2002. Purification and cloning of toxins from elapid venoms that target cyclic nucleotide-gated ion channels. *Biochemistry* 41:11331–37
168. Yamazaki Y, Koike H, Sugiyama Y, Motoyoshi K, Wada T, et al. 2002. Cloning and characterization of novel snake venom proteins that block smooth muscle contraction. *Eur. J. Biochem.* 269:2708–15
169. Yasuda O, Morimoto S, Chen YH, Jiang BB, Kimura T, et al. 1993. Calciseptine binding to a 1,4-dihydropyridine recognition site of the L-type calcium-channel of rat synaptosomal membranes. *Biochem. Biophys. Res. Commun.* 194:587–94
170. Yuan CH, He QY, Peng K, Diao JB, Jiang LP, et al. 2008. Discovery of a distinct superfamily of k-type toxin (KTT) from tarantulas. *PLoS ONE* 3:e3414
171. Zasloff M. 2002. Antimicrobial peptides of multicellular organisms. *Nature* 415:389–95
172. Zhou J, Liao M, Ueda M, Gong H, Xuan X, Fujisaki K. 2007. Sequence characterization and expression patterns of two defensin-like antimicrobial peptides from the tick *Haemaphysalis longicornis*. *Peptides* 28:1304–10
173. Zhou JL, Ueda M, Umemiya R, Battsetseg B, Boldbaatar D, et al. 2006. A secreted cystatin from the tick *Haemaphysalis longicornis* and its distinct expression patterns in relation to innate immunity. *Insect Biochem. Mol. Biol.* 36:527–35

174. Zhu S, Gao B, Tytgat J. 2005. Phylogenetic distribution, functional epitopes and evolution of the CS $\alpha\beta$ superfamily. *Cell. Mol. Life Sci.* 62:2257–69
175. Zhu SY. 2008. Discovery of six families of fungal defensin-like peptides provides insights into origin and evolution of the CS $\alpha\beta$ defensins. *Mol. Immunol.* 45:828–38
176. Zhu SY, Darbon H, Dyason K, Verdonck F, Tytgat J. 2003. Evolutionary origin of inhibitor cysteine knot peptides. *FASEB J.* 17:1765–67
-

Coauthor affiliations:

²Unit of Ecology and Systematics, Vrije Universiteit Brussels, 1050 Brussels, Belgium

³Department of Entomology and Center for Tropical and Emerging Global Diseases, University of Georgia, Athens, Georgia 30602

⁴SBC Lab AG, 8185 Winkel, Switzerland

⁵National School of Pharmacy, University of Otago, Dunedin 9054, New Zealand

⁶Institute for Molecular Bioscience, The University of Queensland, St. Lucia, QLD 4072, Australia

⁷Department of Pathology, University of Turku, Turku, Finland

⁸Sciences Department, Museum Victoria, Melbourne, Victoria 3001, Australia

⁹The Walter and Eliza Hall Institute of Medical Research, Parkville 3050, Victoria, Australia

¹⁰Department of Physiological Sciences, Faculty of Medicine, Pontificia Universidad Javeriana, Bogotá, Colombia

¹¹Structural and Computational Biology/Gene Expression Units, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

**Secondary affiliation: Department of Biochemistry and Molecular Biology, Bio21 Institute, University of Melbourne, Melbourne 3010 Australia



Contents

Chromosomes in Leukemia and Beyond: From Irrelevant to Central Players <i>Janet D. Rowley</i>	1
Unraveling a Multifactorial Late-Onset Disease: From Genetic Susceptibility to Disease Mechanisms for Age-Related Macular Degeneration <i>Anand Swaroop, Emily Y. Chew, Catherine Bowes Rickman, and Gonçalo R. Abecasis</i> ...	19
Syndromes of Telomere Shortening <i>Mary Armanios</i>	45
Chronic Pancreatitis: Genetics and Pathogenesis <i>Jian-Min Chen and Claude Férec</i>	63
The Genetics of Crohn's Disease <i>Johan Van Limbergen, David C. Wilson, and Jack Satsangi</i>	89
Genotyping Technologies for Genetic Research <i>Jiannis Ragousis</i>	117
Applications of New Sequencing Technologies for Transcriptome Analysis <i>Olena Morozova, Martin Hirst, and Marco A. Marra</i>	135
The Posttranslational Processing of Prelamin A and Disease <i>Brandon S. J. Davies, Loren G. Fong, Shao H. Yang, Catherine Coffinier, and Stephen G. Young</i>	153
Genetic Testing in Israel: An Overview <i>Guy Rosner, Serena Rosner, and Avi Orr-Urtreger</i>	175
Stewardship of Human Biospecimens, DNA, Genotype, and Clinical Data in the GWAS Era <i>Stephen J. O'Brien</i>	193
Schistosoma Genetics: New Perspectives on Schistosome Biology and Host-Parasite Interaction <i>Ze-Guang Han, Paul J. Brindley, Sheng-Yue Wang, and Zbu Chen</i>	211
Evolution of Genomic Imprinting: Insights from Marsupials and Monotremes <i>Marilyn B. Renfree, Timothy A. Hore, Geoffrey Shaw, Jennifer A. Marshall Graves, and Andrew J. Pask</i>	241

Methods for Genomic Partitioning <i>Emily H. Turner, Sarah B. Ng, Deborah A. Nickerson, and Jay Shendure</i>	263
Biased Gene Conversion and the Evolution of Mammalian Genomic Landscapes <i>Laurent Duret and Nicolas Galtier</i>	285
Inherited Variation in Gene Expression <i>Daniel A. Skelly, James Ronald, and Joshua M. Akey</i>	313
Genomic Analyses of Sex Chromosome Evolution <i>Melissa A. Wilson and Kateryna D. Makova</i>	333
Sequencing Primate Genomes: What Have We Learned? <i>Tomas Marques-Bonet, Oliver A. Ryder, and Evan E. Eichler</i>	355
Genotype Imputation <i>Yun Li, Cristen Willer, Serena Sanna, and Gonçalo Abecasis</i>	387
Genetics of Athletic Performance <i>Elaine A. Ostrander, Heather J. Huson, and Gary K. Ostrander</i>	407
Genetic Screening for Low-Penetrance Variants in Protein-Coding Diseases <i>Jill Waalen and Ernest Beutler</i>	431
Copy Number Variation in Human Health, Disease, and Evolution <i>Feng Zhang, Wenli Gu, Matthew E. Hurles, and James R. Lupski</i>	451
The Toxicogenomic Multiverse: Convergent Recruitment of Proteins Into Animal Venoms <i>Bryan G. Fry, Kim Roelants, Donald E. Champagne, Holger Scheib, Joel D.A. Tyndall, Glenn F. King, Timo J. Nevalainen, Janette A. Norman, Richard J. Lewis, Raymond S. Norton, Camila Renjifo, and Ricardo C. Rodríguez de la Vega</i>	483
Genetics, Medicine, and the Plain People <i>Kevin A. Strauss and Erik G. Puffenberger</i>	513
Indexes	
Cumulative Index of Contributing Authors, Volumes 1–10	537
Cumulative Index of Chapter Titles, Volumes 1–10	541

Errata

An online log of corrections to *Annual Review of Genomics and Human Genetics* articles may be found at <http://genom.annualreviews.org/>